

The benefits of cardiac rehabilitation for patients with sleep apnea

Saeid Komasi⁽¹⁾ 

Editorial

Date of submission: 23 Mar. 2018, *Date of acceptance:* 12 May 2018

Sleep apnea (SA) is the most common type of breathing-related sleep disorders, defined as a frequent stop and start breathing.¹ SA is characterized by symptoms such as frequent snoring, silent pauses in breathing, choking and breathing symptoms, sleepiness or daily fatigue, insomnia and frequent waking up during the night, morning headaches, not refresh and uneasy feeling after awakening, and irritability and mood changes.² In this disorder, breathing stops occur more than 30 times during sleep, and continue a few seconds to minutes.¹

SA, which is an independent risk factor for cardiovascular disease, affects 38,000 heart deaths each year.^{3,4} SA is common among 50-66 percent of patients with heart diseases, and generally leads to numerous health outcomes.^{5,6} Increased blood pressure due to SA, decreased left ventricular function and central blood flow, myocardial damage, heart rate fluctuations, systemic infection, endothelial dysfunction, increased sympathetic activity, and metabolic abnormalities are among the most important outcome of SA in patients with cardiovascular diseases.^{4,5}

Appropriate treatments for SA include automatic positive airway pressure (APAP) and continuous positive airway pressure (CPAP). Advantages of the APAP method include fluctuation between low and high level pressure during the whole night, and automatically adjust the pressure level, automatic elevation of pressure, and uniform maintenance of pressure level set by patient itself. The limitations of this therapeutic approach are APAP algorithm varies from one person to another; sometimes slowing the pressure change during apnea events, and expensive.¹ The CPAP method, as a non-invasive treatment, quickly relieves symptoms, and alleviates heart problems. An increase in expiratory effort, sense of forced air through the nostrils of the patient, and uniform maintenance of pressure level set by the physician are among the CPAP constraints.¹ This treatment is

generally associated with positive outcomes such as decreased blood pressure caused by SA, and improving the function of the left ventricle and oxygenation.⁵ Despite the usefulness of the APAP and CPAP, SA screening in Iranian patients with cardiovascular diseases is not a routine. For this reason, in most cases, this problem remains underdiagnosed, and without a proper treatment.

Although the aforementioned treatments have been used relatively successful for many years, recent studies have shown that cardiac rehabilitation (CR) can be considered as a new treatment for SA.⁵ CR reduces the severity of SA by 55% through improving VO_2 peak by 20-27 percent.⁷⁻¹⁰ Physical exercise and regular exercise in CR can regulate the autonomic nervous system, and reduce the severity of SA.¹⁰

In spite of sufficient evidence regarding the usefulness of CR in significantly reducing morbidity and mortality,¹¹ Iranian patients tend to have very little to participate in these programs. In Iran, less than 20% of patients [mostly those with coronary artery bypass graft (CABG)] refer to CR, and almost half do not complete the sessions.¹²⁻¹⁴ Not surprisingly, patients with heart failure, valve heart surgery, percutaneous coronary intervention, or myocardial infarction are also not generally encouraged by experts to enroll in CR. Therefore, the number of patients with cardiovascular diseases enrolled in the CR is limited to a number of patients with CABG. Based on these considerations, the numbers of patients with cardiovascular diseases with underdiagnoses/diagnosed SA, who benefits from CR, are very small.

Obviously, SA is a treatable and modifiable disorder.⁴ The above-mentioned literature refers to the benefits of exercise and CR in controlling and reducing the severity of this disorder. CR is effective in controlling and improving SA, though weight loss and body mass index (BMI), modifying the diet, increasing cardiorespiratory fitness,

1 Department of Psychology, Cardiac Rehabilitation Center AND Department of Psychology, Lifestyle Modification Research Center, Kermanshah University of Medical Sciences, Kermanshah, Iran
Correspondence to: Saeid Komasi, Email: s_komasi63@yahoo.com

enhancing VO_2 peak, decreasing leg fluid accumulation, and preventing the nocturnal rostral fluid shift implicated in upper airway collapse.¹⁵ Moreover, other reports refer to the combination of exercise training with ventilation therapy to achieve more positive outcomes.¹⁶ In sum, CR seems to be a potential replacement therapy or complement for existing treatments. However, further studies are needed to confirm this claim.

Acknowledgments

We appreciate the Clinical Research Development Center of Imam Reza Hospital, Kermanshah University of Medical Sciences, Iran.

Conflict of Interests

Authors have no conflict of interests.

References

- Jayaraj R, Mohan J, Kanagasabai A. A review on detection and treatment methods of sleep apnea. *J Clin Diagn Res* 2017; 11(3): VE01-VE03.
- Khazaie H, Najafi F, Rezaie L, Tahmasian M, Sepehry AA, Herth FJ. Prevalence of symptoms and risk of obstructive sleep apnea syndrome in the general population. *Arch Iran Med* 2011; 14(5): 335-8.
- Monahan K, Redline S. Role of obstructive sleep apnea in cardiovascular disease. *Curr Opin Cardiol* 2011; 26(6): 541-7.
- Skobel EC, Kruger S. SERVE-HF: What does it mean for cardiac rehabilitation? *Eur J Prev Cardiol* 2016; 23(2): 125-8.
- Jafari B. Rehabilitation of cardiovascular disorders and sleep apnea. *Sleep Med Clin* 2017; 12(2): 193-203.
- Hargens TA, Aron A, Newsome LJ, Austin JL, Shafer BM. Effects of obstructive sleep apnea on hemodynamic parameters in patients entering cardiac rehabilitation. *J Cardiopulm Rehabil Prev* 2015; 35(3): 181-5.
- Mendelson M, Inami T, Lyons O, Marzolini S, Oh P, Bradley D. Long-term effects of cardiac rehabilitation on sleep apnea severity in patients with coronary artery disease. *Eur Respir J* 2017; 50(Suppl 61): PA4724.
- Mendelson M, Lyons OD, Yadollahi A, Inami T, Oh P, Bradley TD. Effects of exercise training on sleep apnoea in patients with coronary artery disease: A randomised trial. *Eur Respir J* 2016; 48(1): 142-50.
- Mendelson M, Bailly S, Marillier M, Flore P, Borel JC, Vivodtzev I, et al. Obstructive sleep apnea syndrome, objectively measured physical activity and exercise training interventions: A systematic review and meta-analysis. *Front Neurol* 2018; 9: 73.
- Hupin D, Pichot V, Lietar C, Poyraz E, Maudoux D, Sforza E, et al. Impact of cardiac rehabilitation on the obstructive sleep apnoea in the coronary artery disease. *Ann Phys Rehabil Med* 2016; 59: e57.
- Goel K, Lennon RJ, Tilbury RT, Squires RW, Thomas RJ. Impact of cardiac rehabilitation on mortality and cardiovascular events after percutaneous coronary intervention in the community. *Circulation* 2011; 123(21): 2344-52.
- Moradi B, Maleki M, Esmaeilzadeh M, Abkenar HB. Physician-related factors affecting cardiac rehabilitation referral. *J Tehran Heart Cent* 2011; 6(4): 187-92.
- Nalini M. Outpatient cardiac rehabilitation use after coronary bypass surgery in the west of Iran. *J Cardiopulm Rehabil Prev* 2014; 34(4): 263-70.
- Heydarpour B, Saeidi M, Ezzati P, Soroush A, Komasi S. Sociodemographic predictors in failure to complete outpatient cardiac rehabilitation. *Ann Rehabil Med* 2015; 39(6): 863-71.
- Iftikhar IH, Kline CE, Youngstedt SD. Effects of exercise training on sleep apnea: A meta-analysis. *Lung* 2014; 192(1): 175-84.
- Iliou MC, Corone S, Gellen B, Denolle T, Roche F, Bellemain-Apaix A, et al. 0087: Sleep apneas treatment during cardiac rehabilitation can improve heart failure prognosis? SATELIT-HF study: Sleep apnea treatment during cardiac rehabilitation of CHF patients. *Archives of Cardiovascular Diseases Supplements* 2016; 8(1): 30.

How to cite this article: Komasi S. The benefits of cardiac rehabilitation for patients with sleep apnea. *ARYA Atheroscler* 2018; 14(4): 147-8.